Gastroenterology Department

Patient Name:

Your procedure is scheduled for:		at	with Doctor:	
Location:	Gateway Hospital	Deaconess Clinic Downtown	Midtown Hospital	H
	4011 Gateway Blvd	421 Chestnut St	600 Mary St.	1:
	Newburgh, IN 47630	Evansville, IN 47713	Evansville, IN 47710	H

Please check in at the information desk on the first floor, TWO HOURS prior to your appointment time.

If you cannot keep your appointment, please allow a three-day notice for cancellation. To cancel or reschedule your appointment, please call the scheduler at 812-426-9545.

INSTRUCTIONS FOR PREPARING FOR YOUR PROCEDURE

Ticlid

Plavix

**You may receive general anesthesia for this procedure, which can require labs/EKG prior to your procedure. Please call the office as soon as possible at 812-426-9545 to see if any testing is needed.

If you take **blood thinning and/or *weight loss medications, please notify your GI physician as soon as possible at 812-426-9545 for instructions on when/if to stop them. See examples below: Brilinta Xarelto

- Coumadin
- Effient
- Pradaxa
- <u>1 DAY</u> before your procedure
 - 1. Eat a light meal in the evening before your procedure. Stop all food at midnight.

<u>DAY OF</u> your procedure

- 1. **Stop** all water/clear liquids at
- , FOUR hours prior to your procedure time.

Agarenox

Pletal

Arrival Time:

- Stop all candy, mints, chewing tobacco, smoking of all kinds and chewing gum four hours prior to procedure. 2.
- 3. **Stop** diabetic medications unless otherwise directed by your doctor, but bring them with you to the procedure.
- 4. Please take heart, lung, blood pressure and seizure medication with a small sip of water the morning of your procedure.
- Bring your total list of medicines with you to the procedure. 5.
- Bring a responsible adult to receive instructions and provide transportation who will remain at the facility 6. throughout your stay (2-3hrs) You will not be permitted to drive or take public transportation alone (cab, bus, etc) following your exam.
- 7. Wear something comfortable the day of the exam and keep valuables at home.

IMPORTANT: Please follow all instructions listed on this sheet for your safety. Your procedure could be canceled and rescheduled.

- 1. Do not drink clear liquids FOUR hours prior to the procedure. A sip of water with important medicines is okay.
- 2. Stop certain medications prior to procedure (Phentermine for two weeks, blood thinners according to GI MD)
- 3. You must have a responsible adult to receive instructions and provide your transportation the day of your procedure. Taking any public transportation alone (cab, bus, etc) without a responsible adult is not acceptable. This adult must stay with you for the duration of the time you are at the facility (2-3 hrs).
- Follow instructions given by your gastroenterologist or on this instruction sheet ONLY.

Henderson Hospital 1305 N. Elm St. Henderson, KY 42420

Eliquis

- *Phentermine

- And more