

GASTROENTEROLOGY COLONOSCOPY INSTRUCTIONS

Date:	Arrive	e at:	
YOU N	IUST HAVE A DRIVER WITH YOU ON TH	E DAY OF THE PROCEDURE, BECAUSE YOU WILL BE SEDATED.	
before	Hold Coumadin, Brilinta, or Aspirin 5 days prior to procedure once OK'd by prescribing physician.		
SUPPL •		BOTTLE OF MIRALAX POWDER, 64 OZ GATORADE (NO RED OR PURPLE) by pharmacy or Walmart. The Miralax is in a large jug and may be behind is needed.	
DAY B dehyd		uid diet. Drink at least eight glasses of water during the day to avoid	
•	At Noon: Take four Dulcolax tablets. No refrigerator. Do not add ice.	Aix 64 oz. of Gatorade with the Miralax powder. Shake well and place in	
•	At 6:00 p.m.: Drink one 8 oz. glass of to You should have drunk half of the mixt	he Miralax/Gatorade mixture every 20-30 minutes. Stop after four glasses ture.	
-		prior to arrival time to the hospital: Drink an 8 oz. glass of the s until gone. Your stool may appear clear or yellow. This is normal.	
DO NO	T DRINK ANYTHING ELSE		
	· · · · · · · · · · · · · · · · · · ·	or seizure medications with a small sip of water in the morning of the	
colono	scopy. Hold all other medications until a	after the procedure.	
	Clear Liquids	NO RED or PURPLE items	

<u>Clear Liquids</u>	NO RED or PURPLE items
Gatorade, Pedialyte, Powerade	NO alcohol
Clear broth, bouillon	NO mild or non-dairy creamers
Coffee or tea without milk or creamer	NO noodles or vegetables in soup
Carbonated or non-carbonated drinks	NO juice with pulp
Kook-aid, Jell-O, Popsicles, Hard candy	NO liquid you cannot see through
Juice-no pulp	